


Recent Advances in Visualizing Multivariate Linear Models


Michael Friendly Matthew Sigal
with appreciation to Georges Monette & John Fox

May 26–29, 2013, SSC annual Meeting



Outline

- 1 Background
 - Visual overview
 - Data ellipses
 - The Multivariate Linear Model
 - Motivating example
- 2 Hypothesis-Error (HE) plots
 - Visualizing H and E (co)variation
 - MANOVA designs
 - MREG designs
- 3 Reduced-rank displays
 - Low-D displays of high-D data
 - Canonical discriminant HE plots
- 4 Recent extensions
 - Canonical correlation
 - Robust MLMs
 - Influence diagnostics for MLMs
- 5 Conclusions



Slides & R scripts: <http://datavis.ca/papers/ssc2013/>


Introduction: The LM family and friends

Models, graphical methods and opportunities

of response variables

	Classical linear models	Generalized linear models	
1	LM family: $E(y)=X\beta$, $V(y X)=\sigma^2I$ ANOVA, regression, ...	GLM: $E(y)=g^{-1}(X\beta)$, $V=V[g^{-1}(X\beta)]$ poisson, logistic, loglinear, ...	1
	Many graphical methods: effect plots, spread-leverage, influence, ...	Some graphical methods: mosaic plots, 4fold plots, diagnostic plots, ...	
2+	MLM: $E(Y)=X\beta$, $V(Y X)=I\otimes\Sigma$ MANOVA, MMRreg, ...	MGLM: ??? Graphical methods: ???	2
	Graphical methods: ???		


of response variables



Visual overview: Multivariate data, $Y_{n \times p}$

What we know how to do well (almost)


- 2 vars: Scatterplot + annotations (data ellipses, smooths, ...)
- p vars: Scatterplot matrix (all pairs)
- p vars: Reduced-rank display– show max. total variation \mapsto biplot




Visual overview: Multivariate linear model, $Y = XB + U$

What is new here?

- 2 vars: HE plot— data ellipses of H (fitted) and E (residual) SSP matrices
- p vars: HE plot matrix (all pairs)
- p vars: Reduced-rank display— show max. H wrt. $E \mapsto$ Canonical HE plot



Data Ellipses: Galton's data




Galton's data on Parent & Child height: 40%, 68% and 95% data ellipses

Visual overview: Recent extensions

Extending univariate methods to MLMs:

- Robust estimation for MLMs
- Influence measures and diagnostic plots for MLMs
- Visualizing canonical correlation analysis



The Data Ellipse: Details

• Visual summary for bivariate relations

- Shows: means, standard deviations, correlation, regression line(s)
- Defined: set of points whose squared Mahalanobis distance $\leq c^2$,

$$D^2(\mathbf{y}) \equiv (\mathbf{y} - \bar{\mathbf{y}})^T \mathbf{S}^{-1} (\mathbf{y} - \bar{\mathbf{y}}) \leq c^2$$

\mathbf{S} = sample variance-covariance matrix

- Radius: when \mathbf{y} is \approx bivariate normal, $D^2(\mathbf{y})$ has a large-sample χ^2_2 distribution with 2 degrees of freedom.
 - $c^2 = \chi^2_2(0.40) \approx 1$: 1 std. dev univariate ellipse— 1D shadows: $\bar{y} \pm 1s$
 - $c^2 = \chi^2_2(0.68) = 2.28$: 1 std. dev bivariate ellipse
 - $c^2 = \chi^2_2(0.95) \approx 6$: 95% data ellipse, 1D shadows: Scheffé intervals
- Construction: Transform the unit circle, $\mathcal{U} = (\sin \theta, \cos \theta)$,

$$\mathcal{E}_c = \bar{\mathbf{y}} + c\mathbf{S}^{1/2}\mathcal{U}$$

$\mathbf{S}^{1/2}$ = any "square root" of \mathbf{S} (e.g., Cholesky)

- Robustify: Use robust estimate of \mathbf{S} , e.g., MVE (mimimum volume ellipsoid)
- p variables: Extends naturally to p -dimensional ellipsoids

The univariate linear model

- **Model:** $\mathbf{y}_{n \times 1} = \mathbf{X}_{n \times q} \boldsymbol{\beta}_{q \times 1} + \boldsymbol{\epsilon}_{n \times 1}$, with $\boldsymbol{\epsilon} \sim \mathcal{N}(\mathbf{0}, \sigma^2 \mathbf{I}_n)$
- **LS estimates:** $\hat{\boldsymbol{\beta}} = (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}^T \mathbf{y}$
- **General Linear Test:** $H_0: \mathbf{C}_{h \times q} \boldsymbol{\beta}_{q \times 1} = \mathbf{0}$, where \mathbf{C} = matrix of constants; rows specify h linear combinations or contrasts of parameters.
- e.g., Test of $H_0: \beta_1 = \beta_2 = 0$ in model $y_i = \beta_0 + \beta_1 x_{1i} + \beta_2 x_{2i} + \epsilon_i$

$$\mathbf{C}\boldsymbol{\beta} = \begin{bmatrix} 0 & 1 & 0 \\ 0 & 0 & 1 \end{bmatrix} \begin{pmatrix} \beta_0 \\ \beta_1 \\ \beta_2 \end{pmatrix} = \begin{pmatrix} 0 \\ 0 \end{pmatrix}$$

- All \rightarrow F-test: How big is SS_H relative to SS_E ?

$$F = \frac{SS_H/df_h}{SS_E/df_e} = \frac{MS_H}{MS_E} \rightarrow (MS_H - F MS_E) = 0$$

The multivariate linear model

- **Model:** $\mathbf{Y}_{n \times p} = \mathbf{X}_{n \times q} \mathbf{B}_{q \times p} + \mathbf{U}$, for p responses, $\mathbf{Y} = (\mathbf{y}_1, \mathbf{y}_2, \dots, \mathbf{y}_p)$
- **General Linear Test:** $H_0: \mathbf{C}_{h \times q} \mathbf{B}_{q \times p} = \mathbf{0}_{h \times p}$
- Analogs of sums of squares, SS_H and SS_E are $(p \times p)$ matrices, \mathbf{H} and \mathbf{E} ,

$$\mathbf{H} = (\mathbf{C}\hat{\mathbf{B}})^T [\mathbf{C}(\mathbf{X}^T \mathbf{X})^{-1} \mathbf{C}^T]^{-1} (\mathbf{C}\hat{\mathbf{B}}),$$

$$\mathbf{E} = \mathbf{U}^T \mathbf{U} = \mathbf{Y}^T [\mathbf{I} - \mathbf{H}] \mathbf{Y}.$$

- Analog of univariate F is

$$\det(\mathbf{H} - \lambda \mathbf{E}) = 0,$$

- How big is \mathbf{H} relative to \mathbf{E} ?
 - Latent roots $\lambda_1, \lambda_2, \dots, \lambda_s$ measure the "size" of \mathbf{H} relative to \mathbf{E} in $s = \min(p, df_h)$ orthogonal directions.
 - Test statistics (Wilks' Λ , Pillai trace criterion, Hotelling-Lawley trace criterion, Roy's maximum root) all combine info across these dimensions

Motivating Example: Romano-British Pottery

Tubb, Parker & Nicholson analyzed the chemical composition of 26 samples of Romano-British pottery found at four kiln sites in Britain.

- Sites: Ashley Rails, Caldicot, Isle of Thorns, Llanedryn
- Variables: aluminum (Al), iron (Fe), magnesium (Mg), calcium (Ca) and sodium (Na)
- \rightarrow One-way MANOVA design, 4 groups, 5 responses

```
R> library(heplots)
```

```
R> Pottery
```

```

      Site  Al  Fe  Mg  Ca  Na
1  Llanedryn 14.4 7.00 4.30 0.15 0.51
2  Llanedryn 13.8 7.08 3.43 0.12 0.17
3  Llanedryn 14.6 7.09 3.88 0.13 0.20
. . .
25 AshleyRails 14.8 2.74 0.67 0.03 0.05
26 AshleyRails 19.1 1.64 0.60 0.10 0.03
```

Motivating Example: Romano-British Pottery

Questions:

- **Can** the content of Al, Fe, Mg, Ca and Na differentiate the sites?
- **How to understand** the contributions of chemical elements to discrimination?

Numerical answers:

```
R> pottery.mod <- lm(cbind(Al, Fe, Mg, Ca, Na) ~ Site)
R> Manova(pottery.mod)
```

```
Type II MANOVA Tests: Pillai test statistic
      Df test stat approx F num Df den Df Pr(>F)
Site 3      1.55      4.30      15      60 2.4e-05 ***
---
```

```
Signif. codes:  0 '***' 0.001 '**' 0.01 '*' 0.05 '.' 0.1 ' ' 1
```


What have we learned?

- **Can:** YES! We can discriminate sites.
- **But:** **How to understand** the pattern(s) of group differences: ???

Motivating Example: Romano-British Pottery

Univariate plots are limited

- Do not show the *relations* of variables to each other




Motivating Example: Romano-British Pottery

Visual answer: HE plot


- Shows variation of means (**H**) relative to residual (**E**) variation
- Size and orientation of **H** wrt **E**: *how much* and *how* variables contribute to discrimination
- Evidence scaling: **H** is scaled so that it projects outside **E** iff null hypothesis is rejected.

Run heplot-movie.ppt



R> heplot3d(pottery.mod)


HE plots: Visualizing **H** and **E** (co) variation



Ideas behind multivariate tests: (a) Data ellipses; (b) **H** and **E** matrices

- H** ellipse: data ellipse for fitted values, $\hat{y}_{ij} = \bar{y}_j$.
- E** ellipse: data ellipse of residuals, $\hat{y}_{ij} - \bar{y}_j$.


HE plots: Visualizing multivariate hypothesis tests



Ideas behind multivariate tests: latent roots & vectors of **HE**⁻¹

- $\lambda_i, i = 1, \dots, df_h$ show size(s) of **H** relative to **E**.
- latent vectors show canonical directions of maximal difference.

HE plot for iris data



(a) Data ellipses and (b) **H** and **E** matrices (scaled by $1/df_e$: effect size)

- **H** ellipse: data ellipse for fitted values, $\hat{y}_{ij} = \bar{y}_j$.
- **E** ellipse: data ellipse of residuals, $\hat{y}_{ij} - \bar{y}_j$.

HE plot details: **H** and **E** matrices

Recall the data on 5 chemical elements in samples of Romano-British pottery from 4 kiln sites:

R> `summary(Manova(pottery.mod))`

Sum of squares and products for error:

	Al	Fe	Mg	Ca	Na
Al	48.29	7.080	0.608	0.106	0.589
Fe	7.08	10.951	0.527	-0.155	0.067
Mg	0.61	0.527	15.430	0.435	0.028
Ca	0.11	-0.155	0.435	0.051	0.010
Na	0.59	0.067	0.028	0.010	0.199

Term: Site


Sum of squares and products for hypothesis:

	Al	Fe	Mg	Ca	Na
Al	175.6	-149.3	-130.8	-5.89	-5.37
Fe	-149.3	134.2	117.7	4.82	5.33
Mg	-130.8	117.7	103.4	4.21	4.71
Ca	-5.9	4.8	4.2	0.20	0.15
Na	-5.4	5.3	4.7	0.15	0.26

- **E** matrix: Within-group (co)variation of residuals
 - diag: SSE for each variable
 - off-diag: \sim partial correlations
- **H** matrix: Between-group (co)variation of means
 - diag: SSH for each variable
 - off-diag: \sim correlations of means
- How big is **H** relative to **E**?
- Ellipsoids: $\dim(\mathbf{H}) = \text{rank}(\mathbf{H}) = \min(p, df_h)$

HE plot details: Scaling **H** and **E**

- The **E** ellipse is divided by $df_e = (n - p) \rightarrow$ data ellipse of residuals
 - Centered at grand means \rightarrow show factor means in same plot.
- “Effect size” scaling– $\mathbf{H}/df_e \rightarrow$ data ellipse of fitted values.
- “Significance” scaling– **H** ellipse protrudes beyond **E** ellipse iff H_0 can be rejected by Roy maximum root test
 - $H/(\lambda_\alpha df_e)$ where λ_α is critical value of Roy’s statistic at level α .
 - direction of **H** wrt **E** \mapsto linear combinations that depart from H_0 .



R> `heplot(pottery.mod, size="effect")`
 size="evidence")


R> `heplot(pottery.mod,`

HE plot details: Contrasts and linear hypotheses


- An overall effect \mapsto an **H** ellipsoid of $s = \min(p, df_h)$ dimensions
- Linear hypotheses, of the form $H_0 : \mathbf{C}_{h \times q} \mathbf{B}_{q \times p} = \mathbf{0}_{h \times p} \mapsto$ sub-ellipsoid of dimension h , e.g., 2 df test:

$$\mathbf{C} = \begin{bmatrix} 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \end{bmatrix}$$

- 1D tests and contrasts \mapsto degenerate 1D ellipses (**lines**)
- Geometry:
 - Sub-hypotheses are tangent to enclosing hypotheses
 - Orthogonal contrasts form **conjugate axes**



HE plot matrices: All bivariate views



R> pairs(pottery.mod)

AL stands out –
opposite pattern
 $r(\bar{Fe}, \bar{Mg}) \approx 1$


▶ Jump to low-D

HE plots for Multivariate Multiple Regression

- **Model:** $\mathbf{Y} = \mathbf{X}\mathbf{B} + \mathbf{U}$, where cols of \mathbf{X} are quantitative.
- **Overall test:** $H_0 : \mathbf{B} = \mathbf{0}$ (all coefficients for all responses are zero)
 - $\rightarrow \mathbf{C} = \mathbf{I}$ in GLT $\rightarrow \mathbf{H} = \widehat{\mathbf{B}}^T (\mathbf{X}^T \mathbf{X})^{-1} \widehat{\mathbf{B}} = \widehat{\mathbf{Y}}^T \widehat{\mathbf{Y}}$
- **Individual predictors:** $H_0 : \beta_i = 0$
 - $\rightarrow \mathbf{C} = (0, 0, \dots, 1, 0, \dots, 0) \rightarrow \mathbf{H}_i = \widehat{\beta}_i^T (\mathbf{X}^T \mathbf{X})^{-1} \widehat{\beta}_i$
- **HE plot**
 - Overall \mathbf{H} ellipse: how predictors relate collectively to responses
 - Individual \mathbf{H} ellipses ($\text{rank}(\mathbf{H})=1 \rightarrow$ vectors):
 - orientation \rightarrow relation of \mathbf{x}_i to $\mathbf{y}_1, \mathbf{y}_2$
 - length \rightarrow strength of relation
 - collection of individual \mathbf{H} vectors \rightarrow how predictors contribute to overall test.


HE plots for MMRA: Example

- Rohwer data on $n = 37$ low SES children, for 5 PA tasks (N, S, NS, NA, SS) predicting intelligence/achievement (PPVT, SAT, Raven)
- Only NA is individually significant (in this view)
- ... but overall test highly significant
- NA & S contribute to predicting PPVT
- NS & SS contribute to predicting SAT




HE plots for MMRA: MANCOVA

- Rohwer data on $n_1 = 37$ low SES, and $n_2 = 32$ high SES children
- Fit separate regressions for each group
- Are regressions parallel?
- Are they coincident?




HE plots for MMRA: MANCOVA

- Rohwer data on $n_1 = 37$ low SES, and $n_2 = 32$ high SES children
- Fit MANCOVA model (assuming equal slopes)



Low-D displays of high-D data


- High-D data often shown in 2D (or 3D) views— orthogonal projections in variable space— **scatterplot**
- **Dimension-reduction** techniques: project the data into subspace that has the largest *shadow*— e.g., accounts for largest variance.
- → low-D approximation to high-D data



A: minimum-variance projection; B: maximum variance projection


Canonical discriminant HE plots

- As with biplot, we can visualize MLM hypothesis variation for *all* responses by projecting **H** and **E** into low-rank space.
- **Canonical projection:** $\mathbf{Y}_{n \times p} \mapsto \mathbf{Z}_{n \times s} = \mathbf{Y}\mathbf{E}^{-1/2}\mathbf{V}$, where \mathbf{V} = eigenvectors of $\mathbf{H}\mathbf{E}^{-1}$.
- This is the view that maximally discriminates among groups, ie max. **H** wrt **E** !




Canonical discriminant HE plots

- Canonical HE plot is just the HE plot of canonical scores, (z_1, z_2) in 2D, or z_1, z_2, z_3 , in 3D.
- As in biplot, we add vectors to show relations of the y_j response variables to the canonical variates.
- variable vectors here are **structure coefficients** = correlations of variables with canonical scores.




Canonical discriminant HE plots: Properties

- Canonical variates are uncorrelated: **E** ellipse is spherical
- \mapsto axes must be equated to preserve geometry
- Variable vectors show how variables discriminate among groups
- Lengths of variable vectors \sim contribution to discrimination



Canonical discriminant HE plots: Pottery data

- Canonical HE plots provide 2D (3D) visual summary of **H** vs. **E** variation
- Pottery data: $p = 5$ variables, 4 groups $\mapsto df_H = 3$
- Variable vectors: Fe, Mg and Al contribute to distinguishing (Caldicot, Llanedryn) from (Isle Thorns, Ashley Rails): 96.4% of mean variation
- Na and Ca contribute an additional 3.5%. **End of story!**





Run heplot-movie.ppt

Visualizing Canonical Correlation Analysis

- Basic idea: another instance of low-rank approximation

CCA is to *MMReg* as *CDA* is to *MANOVA*



- \rightarrow For quantitative predictors, provides an alternative view of $\mathbf{Y} \sim \mathbf{X}\mathbf{B}$ in space of maximal (canonical) correlations.
- The candisc package implements two new views for CCA:
 - `plot()` method to show canonical (X, Y) variates as **data**
 - `heplot()` method to show **(X, Y)** relations as **heplots** for **Y** in CAN space.



CCA Example: Rohwer data, Ability and PA tests

- `plot()` method shows canonical variates for **X** and **Y** on one dimension
- Smoother shows possible non-linearity
- Point identification highlights unusual observations

```
R> library(candisc)
R> cc <- cancel(cbind(SAT, PPVT, Raven) ~ n + s + ns + na + ss,
+             data=Rohwer, set.names=c("PA", "Ability"))
R> plot(cc, smooth=TRUE, id.n=3)
R> plot(cc, smooth=TRUE, id.n=3, which=2)
```



Robust MLMs

- R has a large collection of packages dealing with robust estimation:
 - `robust::lmrob()`, `MASS::rlm()`, for univariate LMs
 - `robust::glmrob()` for univariate *generalized* LMs
 - High breakdown-bound** methods for robust PCA and robust covariance estimation
 - However, none of these handle the **fully general MLM**
- The `heplots` package now provides `robmlm()` for robust MLMs:
 - Uses a simple M-estimator via iteratively re-weighted LS.
 - Weights: calculated from Mahalanobis squared distances, using a simple robust covariance estimator, `MASS::cov.trob()` and a weight function, $\psi(D^2)$.

$$D^2 = (\mathbf{Y} - \hat{\mathbf{Y}})^T \mathbf{S}_{\text{trob}}^{-1} (\mathbf{Y} - \hat{\mathbf{Y}}) \sim \chi_p^2 \quad (1)$$

- This fully extends the "mlm" class
- Compatible with other mlm extensions: `car::Anova` and `heplots::heplot`.
- Downside: Does not incorporate modern consistency factors or other niceties.

Influence diagnostics for MLMs

- Influence measures and diagnostic plots are well-developed for univariate LMs
 - Influence measures: Cook's D, DFFITS, `dfbetas`, etc.
 - Diagnostic plots: Index plots, `car::influencePlot()` for LMs
 - However, these are have been unavailable for MLMs
- The `mvinfluence` package now provides:
 - Calculation for multivariate analogs of univariate influence measures (following Barrett & Ling, 1992), e.g., Hat values & Cook's D:


$$H_i = \mathbf{X}_i (\mathbf{X}^T \mathbf{X})^{-1} \mathbf{X}_i^T \quad (2)$$

$$D_i = [\text{vec}(\mathbf{B} - \mathbf{B}_{(I)})]^T [\mathbf{S}^{-1} \otimes (\mathbf{X}^T \mathbf{X})] [\text{vec}(\mathbf{B} - \mathbf{B}_{(I)})] \quad (3)$$

- Provides deletion diagnostics for subsets (I) of size $m \geq 1$.
- e.g., $m = 2$ can reveal cases of **masking** or **joint influence**.
- Extension of `influencePlot()` to the multivariate case.
- A new plot format: **leverage-residual (LR) plots** (McCulloch & Meeter, 1983)

Robust MLMs: Example


For the Pottery data:



- Some observations are given weights ~ 0
- The **E** ellipse is considerably reduced, enhancing apparent significance

Influence diagnostics for MLMs: Example

For the Rohwer data:




Cook's D vs. generalized Hat value

Leverage - Residual (LR) plot

Influence diagnostics for MLMs: LR plots

- Main idea: Influence \sim Leverage (L) \times Residual (R)
- $\mapsto \log(\text{Infl}) = \log(L) + \log(R)$
- \mapsto contours of constant influence lie on lines with slope = -1.
- Bubble size \sim influence (Cook's D)
- This simplifies interpretation of influence measures



Conclusions: Graphical methods for MLMs

Summary & Opportunities

- **Data ellipse:** visual summary of bivariate relations
 - Useful for multiple-group, MANOVA data
 - Embed in scatterplot matrix: pairwise, bivariate relations
 - Easily extend to show partial relations, robust estimators, etc.
- **HE plots:** visual summary of multivariate tests for MANOVA and MMRA
 - Group means (MANOVA) or 1-df H vectors (MMRA) aid interpretation
 - Embed in HE plot matrix: all pairwise, bivariate relations
 - Extend to show partial relations: HE plot of "adjusted responses"
- **Dimension-reduction techniques:** low-rank (2D) visual summaries
 - Biplot: Observations, group means, biplot data ellipses, variable vectors
 - Canonical HE plots: Similar, but for dimensions of maximal discrimination
- **Beautiful and useful geometries:**
 - Ellipses everywhere; eigenvector-ellipse geometries!
 - Visual representation of significance in MLM
 - Opportunities for other extensions

— FIN —